

PHYS 106 Section 001 Problem Set 1 Answer Key

① i) Formula for kinetic energy = $E = \frac{1}{2}mv^2$

$$1 \text{ J} = \frac{1 \text{ kg} \cdot \text{m}^2}{\text{s}^2}$$

From the problem:

$$25 \text{ km/s} = \text{velocity } (v)$$

$$2.5 \times 10^5 \text{ kg} = \text{mass of spaceship}$$

$$E = \left(\frac{1}{2}\right) (2.5 \times 10^5 \text{ kg}) \left(\frac{25 \text{ km}}{\text{s}} \cdot \frac{1000 \text{ m}}{\text{km}}\right)^2$$

$$E = \left(\frac{1}{2}\right) (2.5 \times 10^5 \text{ kg}) (6.25 \times 10^8 \text{ m}^2/\text{s}^2)$$

$$E = 7.81 \times 10^{13} \text{ J}$$

ii) Radius of nucleus = $1.5 \times 10^{-5} \text{ pm} \times \frac{1 \times 10^{-12} \text{ m}}{\text{pm}} = 1.5 \times 10^{-17} \text{ m}$

iii) Mass of carbon atom = $2.0 \times 10^{-23} \text{ g}$

Density of carbon nucleus = $\frac{\text{mass of carbon nucleus}}{\text{volume of carbon nucleus}}$

Volume of carbon nucleus = $\frac{4}{3} \pi (r^3) = 1.51 \times 10^{-50} \text{ m}^3$

$$\text{Density of Carbon nucleus} = \frac{1.32 \times 10^{27} \text{ g}}{\text{m}^3}$$

b) Radius of earth if density of earth equalled
Cerbera nucleus density

$$\text{Mass of earth} = 5.97 \times 10^{24} \text{ kg} \times \frac{1000 \text{ g}}{\text{kg}} = \boxed{5.97 \times 10^{27} \text{ g}}$$

$$\text{Density} = \frac{1.32 \times 10^{27} \text{ g}}{\text{m}^3} = \frac{5.97 \times 10^{27} \text{ g}}{\text{m}^3}$$

$$x = \frac{5.97 \times 10^{27} \text{ g}}{1.32 \times 10^{27} \text{ g/m}^3} = 4.52 \text{ m}^3 = V$$

$$x = 4.52 \text{ m}^3 = \frac{4}{3} \pi r^3 \text{ (volume of earth if)}$$

$$r^3 = 7.1077 \text{ m}^3 \quad \text{density of earth = density of Cerbera nucleus}$$

$$\boxed{r = 1.02 \text{ m}}$$

Density is in g/cm^3 , so:
mass of the earth is:


$$5.97 \times 10^{24} \text{ kg} \times \frac{1000 \text{ g}}{\text{kg}} = 5.97 \times 10^{27} \text{ g}$$

Therefore, the density would be:

$$\text{Density } D = \frac{\text{mass}}{\text{volume}} = \frac{5.97 \times 10^{27}}{1.11 \times 10^{11} \text{ cm}^3} = \boxed{4.23 \times 10^7 \text{ g}/\text{cm}^3}$$

iii) In this problem, we are writing a base with its conjugate acid and letting them sit @ equilibrium.

This problem could be done using the K_B for the reaction



$$K_B = \frac{[\text{NH}_4^{\oplus}][\text{OH}^{\ominus}]}{[\text{NH}_3]}$$

$$[\text{OH}^{\ominus}] = \frac{K_B [\text{NH}_3]}{[\text{NH}_4^{\oplus}]}$$

$$[\text{OH}^{\ominus}] = \frac{(1.8 \times 10^{-5})(0.125 \text{ M})}{(0.3 \text{ M})}$$


$$\boxed{[\text{OH}^{\ominus}] = 7.5 \times 10^{-6}}$$

We know the molarity of NH_3 and NH_4^{\oplus} so plug and chug after rearranging to solve for $[\text{OH}^{\ominus}]$

iv) The titration of methylamine with H_3O^+ has a chemical reaction that looks like:



At the equivalence point, all of the methylamine has been converted to methyl ammonium, so the major species is



2) Use the graph! The equivalence point is reached when 10ml of base has been added and the pH is $\boxed{8.0}$

i) If the equivalence point is reached @ 10ml, then 25% of the equivalence point would be $0.25 \times 10\text{ml} = 2.5\text{ml}$. The pH @ 2.5ml of base added is $\sim \boxed{3.5}$

iii) Halfway to the equivalence point, the pH would be $\boxed{14}$

iv) 75% of the way to the equivalence point, the volume of base added would be 7.5ml and the pH is $\boxed{14.5}$

- ③ Both the acid and the base are the same concentration. We had 50ml of acid and added 55ml of base. We have neutralized all of the acid and have 5ml of OH left over.

New total volume: 105ml (volume of acid + volume of base)
 moles of base in excess: $0.005 \cancel{L} \times \frac{0.02 \cancel{mol}}{\cancel{L}} = 1 \times 10^{-4} \text{ moles OH}^-$

$$pH + pOH = 14$$


$$pH = 14 - pOH$$

$$pOH = -\log [OH^-] = -\log \left(\frac{1 \times 10^{-4} \text{ moles OH}^-}{0.105 \text{ L}} \right)$$

$$pOH = 3.02$$

$$pH = 14 - 3.02 = \boxed{10.98}$$

- ④ From the problem, you can see that there are 2 acid protons



Just like problem 2, the first endpoint is @ the steepest slope, so 50ml of OH added.

Halfway to this volume is pKa1, so when 25ml of OH has been added, the pH is $\boxed{2}$

using the same logic for the second proton, the endpoint volume is 100ml of OH added, so the halfway point would be 75ml of OH added. the pH is $\boxed{7}$

5) i) If the interaction energy is proportional to $1/r^2$ we know we have an ion-dipole interaction.

Chloromethane isn't an ion, so (a) is out. Na^+ is an ion and H_2O is a dipole. This is the correct answer.

B

ii) An interaction energy proportional to $1/r^3$ is a dipole-dipole interaction. Find 2 dipoles or polar molecules.

Chloromethane is polar, and would interact in the liquid phase.

Choice A is correct

Ionic solid isn't a dipole so (B) is incorrect.

Bromine molecules (Br_2) isn't polar so (C) is incorrect.

Chloromethane in the solid phase has some peculiarities that you are unaware of, and some chloromethane is polar, **Choice D is correct**.

iii) We are looking for Dipole/Induced dipole or London Force interactions.

Chloromethane is polar so (A) is incorrect. So is (B).

Li^+ is an ion and H_2O is a dipole so (C) is incorrect.

Na^+ and H_2O , same thing.

This doesn't have an answer that makes sense according to what we talked about in class.

Free points!!! yay!

iv) Highest molar mass = highest boiling point
since all but choice (C) are polar

Answer = (B)

v) Hydrogen bonding is the strongest interaction