

# Answer Key


PHYS 201 Fall 2019 Test #3 Name: \_\_\_\_\_

I. For the following multiple choice questions, write your answer in the line next to the question number.

1-2) Moon takes 27 days to make 1 rotation about its own axis.

1. What is the angular speed of moon's rotation in degree/day?  $\frac{360}{27}$
- a. 6      **(b.) 13**      c. 15      d. 30      e. 36       $\frac{2\pi}{27 \text{ days}} \times \frac{1 \text{ day}}{24 \text{ hrs}} \times \frac{1 \text{ hr}}{60 \text{ min}} \times \frac{1 \text{ min}}{60 \text{ s}}$
2. What is the angular speed of moon's rotation in rad/s?
- a. 0.105      b. 0.556      c.  $1.75 \times 10^{-3}$       d.  $9.7 \times 10^{-3}$       **(e.)  $2.7 \times 10^{-6}$**       f.  $1.6 \times 10^{-4}$

3. The drawing illustrates an overhead view of a door and its axis of rotation. The axis is perpendicular to the page. There are four forces acting on the door, and they have the same magnitude. Which force will provide the highest torque, about the axis of rotation?



- a.  $F_1$   
b.  $F_2$   
**(c.)  $F_3$**   
d.  $F_4$


longer  
LA

4-6) A uniform meter stick is supported at the 25 cm mark using a 12-g knife-edge clamp. Balance is obtained when a 55 gram mass is suspended at the 10 cm mark.

4. Draw a free-body diagram for the meter stick.

5. What is the mass of the meter stick?

6. What is the normal force in gram?



$33 + 12 + 55 = 100$

Answers for 5 & 6

- a. 22 g      **(b.) 33 g**      c. 75 g  
d. 88 g      e. 93 g      f. 100 g

7. When a 54 kg man stands on a pogo stick, the spring is compressed 18 cm. What is the spring constant of the spring, in SI units? ( $g = 9.8 \text{ m/s}^2$ )

- a. 3.0      b. 29.4      c. 300      **(d.) 2940**      e. 29400

8-10) A point mass, 2 kg is located at (3m, 4m) in the coordinate system below. Z axis is perpendicular to the X and Y axes.


8. What is the moment of inertia of this mass about the x axis?

9. What is the moment of inertia of this mass about the y axis?

10. What is the moment of inertia of this mass about the z axis?

Answers for 8-10

- a.  $18 \text{ kg}\cdot\text{m}^2$       b.  $32 \text{ kg}\cdot\text{m}^2$       c.  $36 \text{ kg}\cdot\text{m}^2$   
d.  $50 \text{ kg}\cdot\text{m}^2$       e.  $64 \text{ kg}\cdot\text{m}^2$       f.  $100 \text{ kg}\cdot\text{m}^2$



$I = mr^2$

$I_x = (2)(4^2) = 2 \times 16 = 32$


$I_y = (2)(3^2) = 2 \times 9 = 18$

$I_z = (2)(5^2) = 2 \times 25 = 50$

$55(15) = (m)(25)$   
 $825 = 25m$   
 $m = 33$

$54(9.8) = k(0.18)$   
 $529.2 = 0.18k$   
 $2940$

- C 11. Sit-ups are more difficult to do with your hands placed behind your head instead of on your stomach. This is because,
- The mass is greater when the hands are placed behind the head instead on the stomach.
  - The mass is smaller when the hands are placed behind the head instead on the stomach.
  - The moment of inertia is greater when the hands are placed behind the head instead on the stomach.
  - The moment of inertia is smaller when the hands are placed behind the head instead on the stomach.



A 12. What is the amplitude, in m, for the wave shown above?

E 13. What is the period, in s, for the wave shown above?

B 14. What is the frequency in Hz for the wave shown above?

Answers for 12-14:

- a. 0.20      b. 0.25      c. 0.40      d. 0.50      e. 4.0

$$f = \frac{1}{T} = \frac{1}{4}$$

A 15. Pendulum clocks are made to run at the correct rate by adjusting the pendulum's length. Suppose you move from one city to another where the acceleration due to gravity is slightly greater, taking your pendulum clock with you, what will you have to do to the length of the pendulum to keep the correct time, other factors remaining constant?

Period of a simple pendulum:  $T = 2\pi \sqrt{\frac{L}{g}}$

$g \uparrow \quad L \uparrow$

- a. lengthen it      b. shorten it      c. keeping the length same

End of MC questions

II. A playground merry-go-round with a child at the rim is rotating with an angular velocity of 2.5 rad/s. If the mass of the child is 19 kg, what will be the angular velocity of the merry-go-round as soon as the child jumps out along a radial direction? (Assume that the merry-go-round is a disk of mass 110 kg and radius 2.2 m)




$$I_1 \omega_1 = I_2 \omega_2$$

merry-go-round + kid  
 $(358.16) \frac{2.5 \text{ rad}}{\text{s}} = 266.2 (\omega_2)$

$$895.4 = 266.2 \omega_2$$

$$\omega_2 = \frac{3.36 \text{ rad}}{\text{s}}$$



$$I = \frac{1}{2} MR^2 \text{ merry-go-round}$$

$$I = \frac{1}{2} (110)(2.2^2)$$


$$266.2$$

$$I = mR^2 \text{ kid}$$

$$19(2.2^2)$$

$$91.96$$

III. A pulley of moment of inertia  $8.5 \text{ kg}\cdot\text{m}^2$  and radius  $0.15 \text{ m}$  is pulled with an applied force,  $F = 18 \text{ N}$ , as shown below.



1. Calculate the torque exerted on the pulley.

$$T = LA \cdot F$$

$$T = (0.15)(18) = 2.7 \text{ Nm} = T$$

2. Calculate the angular acceleration of the pulley.

$$T = I\alpha$$

$$2.7 = 8.5\alpha$$

$$\alpha = 0.32 \text{ rad/s}^2$$

3. If the pulley is at rest initially, how long the force needs to be applied to make the pulley spin at  $26 \text{ rad/s}$ ?  $w = \omega$   $w_0 = 0$   $\omega = ?$

$$\omega = \omega_0 + \alpha t$$

$$26 = 0 + 0.32t$$

$$t = 82 \text{ s}$$

IV. Period of oscillating mass on spring:

$$T = 2\pi \sqrt{\frac{m}{k}}$$

1. Show that the above equation is correct unit wise.

$$T = 2\pi \sqrt{\frac{\text{kg}}{\text{N/m}}}$$

$$S = 2\pi \sqrt{\frac{\frac{\text{kg}}{\text{m}}}{\frac{\text{kg}\cdot\text{m}}{\text{s}^2}}}$$

$$S = 2\pi \sqrt{\text{s}^2}$$

$$S = S$$

$$N = \frac{\text{kg}\cdot\text{m}}{\text{s}^2}$$

$$\frac{\text{kg}\cdot\text{s}^2}{\text{kg}\cdot\text{m}} \cdot \frac{\text{m}}{1} = \text{s}^2$$

2. A spring-mounted chair in which the astronaut sits, can be used to find the mass of an astronaut. The chair is then made to oscillate in simple harmonic motion. The spring used in one such device has a spring constant of  $596 \text{ N/m}$ , and the mass of the chair is  $13.0 \text{ kg}$ . The measured oscillation period is  $2.21 \text{ s}$ . Find the mass of the astronaut.

$$T = 2\pi \sqrt{\frac{m}{k}}$$

$$2.21 = 2\pi \sqrt{\frac{m}{596}}$$

$$\frac{2.21}{2\pi} = \sqrt{\frac{m}{596}}$$

$$\left(\frac{2.21}{2\pi}\right)^2 \cdot 596 = m$$

$$m = 73.7 - 13 = 60.7 \text{ kg for astronaut}$$

V. The spring-loaded gun in a ballistic pendulum is compressed by 4.3 cm and a 49 gram steel ball is released. If the spring constant of the spring is 750 N/m, calculate the speed of the ball as it leaves.

$$\frac{1}{2} kx^2 = \frac{1}{2} mv^2$$


$$\frac{1}{2} (750) (0.043)^2 = \frac{1}{2} (0.049) (v)^2$$

$$0.693 = \frac{1}{2} (0.049) (v)^2$$

$$28.3 = v^2$$

$$v = 5.31 \text{ m/s}$$

VI. A 1500-N uniform horizontal beam of length 8.0 m is hinged to a vertical wall at one end and is supported by a cable at the other end as shown below. A fireman of mass 830-N is standing on the beam at 5.0 m from the hinge. Draw a free-body diagram for the beam, showing and naming all the forces acting on the beam, including the horizontal ( $R_x$ ) and vertical ( $R_y$ ) forces exerted by the hinge on the beam.



$$R_x = (1500 \times 4) + (830 \times 5)$$

$$R_y - \text{Tension} = 0 \quad R_y = \text{Tension} \cos$$

VII. A man holds a 165-N ball in his hand, with the forearm horizontal (see the drawing below). He can support the ball in this position because of the flexor muscle force,  $\vec{M}$ , which is applied perpendicular to the forearm and the force,  $\vec{B}$ , applied by the upper-arm bone.

The forearm weighs 21.0 N and has a center of gravity as indicated.

a) Show the weight of the ball and forearm in the diagram.


b) Find the magnitudes of the forces  $\vec{M}$  and  $\vec{B}$ .

$$(33 \times 165) + (21 \times 14) + (M \times 5) = 0$$

$$5445 + 294 = 5M$$

$$5739 = 5M$$

$$1147.8 \text{ N} = \vec{M}$$



Arm not moving therefore forces in each direction (up and down) must be equal

$$\text{Down} \downarrow \text{ forces} = \vec{B} + 21 \text{ N} + 165 \text{ N} = 1148 \text{ N} \quad \text{Up} \uparrow \text{ forces}$$

$$\vec{B} = 962 \text{ N}$$

$$\vec{B} \perp \vec{M} = 0$$

$$F = ? \times$$